

Oral Health Fact Sheet for Medical Professionals

Children with Anxiety and Depression

Anxiety is characterized by apprehension or fear of impending actual or imagined danger, vulnerability, or uncertainty and may be accompanied by restlessness, tension, tachycardia, and dyspnea unattached to a clearly identifiable stimulus. (ICD 9 code 300.0)

Depression is an unpleasant, but not necessarily irrational or pathological, mood state characterized by sadness, despair, or discouragement; it may also involve low self-esteem, social withdrawal, and somatic symptoms such as eating and sleep disturbance. (ICD 9 code 311.0)

Oral Manifestations and Considerations

Oral

- Neglect of oral hygiene leading to increased risk of dental caries and periodontal disease
- Poor nutrition
- Drug-induced xerostomia
- Some studies report a positive correlation with temporomandibular disorders and atypical facial pain.

Other Potential Disorders/Concerns

- Mitral valve prolapse and GERD
- Children with depression are at increased risk for engaging in high-risk behaviors (promiscuity, smoking, alcohol and drug abuse)

Oral Side Effects of Commonly Prescribed Medications

Medication

SYMPTOM	MEDICATION	SIDE EFFECTS
Depression	SSRIs, Atypical Antidepressants, and Tricyclic Antidepressants (TCA's)	Xerostomia, dysphagia, sialadenitis, dysgeusia, stomatitis, gingivitis, glossitis, tongue edema, discolored tongue, and bruxism.
Anxiety	SSRIs, Atypical Antidepressants, and Benzodiazepines	Xerostomia, dysphagia, sialadenitis, dysgeusia, stomatitis, gingivitis, glossitis, tongue edema, discolored tongue, and bruxism.

Children with Anxiety and Depression continued

Parent/Caregiver Support and Guidance

- Discourage consumption of cariogenic foods and beverages
- Prescribe sugar-free medications if available
- Recommend preventive measures such as topical fluoride and sealants
- Advise the use of fluoridated toothpaste twice daily and support the family in following dental care instructions
- Instruct caregiver on appropriate protocol following dental trauma (seek immediate professional care and locate/preserve missing tooth)
- Recommend rinsing with water thoroughly after taking each dose of sugar containing medication and frequent water intake for children taking xerostomic medications
- Recommend preventive measures such as topical fluoride and sealants
- Review safety issues appropriate to the age of the child, such as mouth guards to prevent oral-facial trauma
- Refer to dentist any oral developmental abnormalities

Further information (Medical Anticipatory Guidance, Dental Anticipatory Guidance, Oral Health Guidance for parents/Caregivers and Dental Professional Fact Sheet for Anxiety and Depression) can be found at:

http://dental.washington.edu/departments/omed/decod/special_needs_facts.php

References

- Friedlander, A.H., Mahler, M.E. (2001) Major depressive disorder: Psychopathology, medical management and dental implications. J Am Dent Assoc, 132(5): 629-638.
- Becker, D. E. (2008) Psychotropic drugs: implications for dental practice. Anesthesia Progress, 55(3): 89-99.
- Deykin, EY., Buka, S.L., Zeena, T.H. (1992) Depressive illness among chemically dependent adolescents. Am J Psychiatry, 149(10):1341-1347.

Additional Resources

- [NIH Institute for Depression Disorder](#) and [NIH Institute for Anxiety Disorder](#)
- [Special Care: an Oral Health Professionals Guide to Serving Young Children with Special Health Care Needs](#)
- [Bright Futures Oral Health Pocket Guide](#)
- [American Academy of Pediatrics Oral Health Initiative](#)
- [American Academy of Pediatric Dentistry: 2008-09 Definitions, Oral Health Policies and Clinical Guidelines](#)
- [MCH Resource Center](#)
- [ASTDD-Special Needs](#)
- [Block Oral Disease, MA](#)
- [NOHIC-NIDCR publications](#)



Dear Colleague,

The Washington State Department of Health (DOH) provides print-ready files (PDFs) of health education materials. To ensure that the original quality of the piece is maintained, please read and follow the instructions below and the specifications included for professional printing.

- **Use the latest version.** DOH materials are developed using the most current information available, are checked for clinical accuracy, and are field tested with the intended audience to ensure they are clear and readable. DOH programs make periodic revisions to educational materials, so please check this web site to be sure you have the latest version. DOH assumes no responsibility for the use of this material or for any errors or omissions.
- **Do not alter.** We are providing this artwork with the understanding that it will be printed without alterations and copies will be free to the public. Do not edit the text or use illustrations or photographs for other purposes without first contacting us. Please do not alter or remove the DOH logo, publication number or revision date. If you want to use a part of this publication for other purposes, contact the Office of Health Promotion first.
- **For quality reproduction:** Low resolution PDF files are intended for black and white or color desktop printers. They work best if you are making only one or two copies. High resolution PDF files are intended for reproducing large quantities and are set up for use by professional offset print shops. The high resolution files also include detailed printing specifications. Please match them as closely as possible and insist on the best possible quality for all reproductions.

If you have questions, contact:
Office of Health Promotion
P.O. Box 47833 Olympia, WA 98504-7833
(360) 236-3736

Sincerely,
Health Education Resource Exchange Web Team